


AGUIDE TO THE ELEMENTS AGUIDE TO THE

SECOND EDITION



Гhe Periodic Table		6
Hydrogen16	Cerium	148
Helium22	Praseodymium	
Lithium25	Neodymium	
Beryllium29	Promethium	
Boron31	Samarium	
Carbon34	Europium	
Nitrogen41	Gadolinium	
Oxygen47	Terbium	159
Fluorine50	Dysprosium	160
Neon53	Holmium	
Sodium54	Erbium	
Magnesium57	Thulium	
Aluminum59	Ytterbium	164
Silicon61	Lutetium	
Phosphorus64	Hafnium	
Sulfur66	Tantalum	
Chlorine69	Tungsten	
Argon71	Rhenium	
Potassium73	Osmium	
Calcium75	Iridium	
Scandium79	Platinum	
Titanium81	Gold	
Vanadium83	Mercury	
Chromium85	Thallium	
Manganese87	Lead	
Iron89	Bismuth	
Cobalt92	Polonium	
Nickel94	Astatine	
Copper96	Radon	
Zinc99	Francium	
Gallium101	Radium	
Germanium103	Actinium	
Arsenic104	Thorium	
Selenium104	Protactinium	
Bromine108	Uranium	
Krypton110	Neptunium	
Rubidium112	Plutonium	207
Strontium113	Americium	
Yttrium115	Curium	
Zirconium117	Berkelium	
Niobium119	Californium	
Molybdenum121	Einsteinium	
Technetium122	Fermium	
Ruthenium124	Mendelevium	
Rhodium125	Nobelium	
Palladium126	Lawrencium	
Silver127	Rutherfordium	
Cadmium130	Dubnium	
Indium132	Seaborgium	
Tin	Bohrium	
Antimony135	Hassium	
Tellurium136	Meitnerium	
Iodine137	Ununnilium	
Xenon140	Unununium	
Cesium142	Ununbiium	
Barium144	Ununquadium	
Lanthanum146	Epilogue	
Editification 140	2720840	
		2.2
Glossary		
Chronology		.237
Further Reading		.241
Web Sites		
Index		.243

OXFORD

A Guide to the ELEMENTS

SECOND EDITION

ALBERT STWERTKA

OXFORD UNIVERSITY PRESS

NEW YORK • OXFORD

Oxford University Press

Oxford New York Auckland Bangkok Buenos Aires Cape Town Chennai Dar es Salaam Delhi Hong Kong Istanbul Karachi Kolkata Kuala Lumpur Madrid Melbourne Mexico City Mumbai Nairobi São Paulo Shanghai Singapore Taipei Tokyo Toronto

and an associated company in

Berlin

Copyright © 1996, 2002 by Albert Stwertka Published by Oxford University Press, Inc. 198 Madison Avenue, New York, New York 10016

Oxford is a registered trademark of Oxford University Press.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior permission of Oxford University Press.

Consultant: Robin Eichen Conn, Cargill, Inc.

Design: Valerie Sauers

Picture research: Amla Sanghvi, Jennifer Traslavina

Library of Congress Cataloging-in-Publication Data Stwertka, Albert

A guide to the elements / Albert Stwertka. — 2nd ed.

p. cm.

Includes bibliographical references and index.

1. Chemical elements—Juvenile literature. 2. Periodic law—Juvenile literature. [1. Chemical elements.] I. Title.

QD466.S78 2002 546' .8—dc21 98-20372

C

CIP

AC

ISBN 0-19-515026-0 (lib. ed.); ISBN 0-19-515027-9 (pb. ed.)

98765432

Printed in Hong Kong on acid-free paper

The Periodic Table	6
Hydrogen16	Cerium148
Helium22	Praseodymium149
Lithium25	Neodymium151
Beryllium29	Promethium153
Boron31	Samarium155
Carbon34	Europium156
Nitrogen41	Gadolinium157
Oxygen47	Terbium159
Fluorine50	Dysprosium160
Neon53	Holmium161
Sodium54	Erbium162
Magnesium57	Thulium163
Aluminum59	Ytterbium164
Silicon61	Lutetium165
Phosphorus64	Hafnium166 Tantalum168
Sulfur	
Chlorine	Tungsten
Argon71 Potassium73	Osmium
Calcium	Iridium174
Scandium79	Platinum175
Titanium81	Gold
Vanadium83	Mercury181
Chromium85	Thallium184
Manganese87	Lead186
Iron89	Bismuth189
Cobalt92	Polonium191
Nickel94	Astatine193
Copper96	Radon194
Zinc99	Francium196
Gallium101	Radium197
Germanium103	Actinium199
Arsenic104	Thorium200
Selenium106	Protactinium202
Bromine108	Uranium203
Krypton110	Neptunium206
Rubidium112	Plutonium207 Americium209
Strontium113	Curium210
Yttrium115 Zirconium117	Berkelium211
Niobium119	Californium212
Molybdenum121	Einsteinium214
Technetium	Fermium215
Ruthenium124	Mendelevium216
Rhodium125	Nobelium217
Palladium126	Lawrencium218
Silver127	Rutherfordium219
Cadmium130	Dubnium221
Indium132	Seaborgium222
Tin133	Bohrium224
Antimony135	Hassium225
Tellurium136	Meitnerium226
Iodine137	Ununnilium227
Xenon	Unununium
Cesium142	Ununbiium
Barium	Ununquadium230
Lanthanum146	Epilogue232
Glossary	234
Chronology	237
Further Reading	241
	242
Index	243

1	ĺ		į		ľ	l				ļ			ŀ	ĺ								
	į	ĺ		(1													١		į		
1		Ì		ĺ										ĺ	ĺ							
	į			į											1				2			
		í	۱		Ì)										ĺ)	
					ı			į	I				ĺ								-	

			\int_{∞}			<u> </u>		\sum_{\sim}		* `	<u> </u>		
18 VIII.A	10 Z 20.1797	≱	39.948	% _	83.80	54	Xe	131.29	98	8	110.777		
VIIA	9 18.99840	ะบ	35.4527	55 Q	79.904	53	****	126.09447	85	At* *	207.707		
91 VIA	8 O 15.9994	့ ဟ	32.066	% Se	78.96	52	౼	121.757	84	Po **	200.7024		
15 VA	7 Z	<u> </u>	30.97376	33 A S	74.92159	51	Sp	121.76	83	. <u> </u>			_
4 NA	و ا2.011	± ஜ	28.0855	% G	72.61	50	Sn	118.710	82	2		**DUQ	1300
13 L3	5 B 10.811	A	26.98154	~ B	69.723	49		114.82	81	—	201-2020		_
`		2 =		္က ဂ	62.39	48	ಕ	112.411	80	ာ	112	Lub*	1770
	genera	<u>8</u>		~ ∂	63.546	47	Ag	107.8682	79	Ac Second		**nnn	1070
Atomic number Chemical symbol Atomic weight	2	2		₹	58.6934	46	Pd	106.42	78	⊈ 105 08	011	*	
Atomic number Chemical symbo Atomic weight	9 A				58.93320	45	된	102.90550	77	7 201	109	*	
Y Y Y	α			ှ မှ	55.847	44		101.07	9/	ဝိန္	108	*	
- T		VIIB	1	٩ ₹	54.9380	43	* ₂	28.9072	75	Se 2012	107	*	
Company and Compan	~	VIB		-	1966.15	42	Š	ᅱ	74	3 83.85	106	*	o i
	Ľ	NB N		?>	50.9415	41	<u>S</u>	92.90638	7.3	D	105	*	262 114
	4	IVB	1	7 =	47.88	40		91.224 (72		+	*	76111
process	r	IIIB		⊸ သိ	44.95591	39	>	88.90585	57	138 9055	68	*	9770776
2 IIA	901218	S	24.3050	⊲	_	38		87.62	56		88	*	. 1250,450
■ - ■ - ■ +62000	6.941	2			39.0983	37	2	85.4678	55	CS 39 90543	87	*	773 0197
	7	Ω,		boin 4			١٠			9			

	-			*	<u>-</u>
17	32.5			*	
	<u>ڳ</u>				
69	E			* Wd*	
89			3	* * *	257.0951
29	2	104.75032	66	ES*	252.083
99	<u>ک</u>			ţ,	251.0796
65	4	130.72337	/6	**	247.0703
64	8	67.76	76	* E	247.0703
63	3 2 131	101.700	۲۶	Am*	243.0614
29	E _S	00.00	74	* _	244.0642
19	Pa **	771.7127	73	* 2	237.0482
09	Z 2	F 7.T.F.	76	* 5	238.0289
59	4 00 90765	0.00.00	7	* * *	223.0359
58	9 5	011.05	2	<u>*</u>	222.0381
	*		-		•

**All the isotopes of this element are radioactive. With the exception of uranium and thorium, the atomic weight shown represents the relative atomic weight of the longest-lived isotope. The numbers in parentheses are the mass numbers of the longest-lived isotope.

Ithough the world is varied and complex, everything in it—air, water, rocks, living tissue, and the almost infinite number of other objects and materials around us—is actually made up of only a limited number of chemical elements. We know today that only 91 such elements exist naturally on the Earth. They range from hydrogen, the lightest element, to uranium, the heaviest. Actually, several more elements do exist, but these have to be made artificially in laboratories.

The basic components of each chemical element are atoms. The atoms of an element consist of three kinds of particles: protons, neutrons, and electrons. Protons and neutrons exist at the core, or nucleus, of the atom. One of the important ways in which these two kinds of particles differ from one another is that each proton carries a single, positive electric charge, whereas a neutron carries no electric charge. Electrons, which are much smaller than either protons or neutrons, each carry a single negative electric charge. Electrons are present at some distance away from the nucleus of the atom and travel rapidly around it in complex paths known as orbits. Under normal circumstances, the number of electrons orbiting around the nucleus of a particular atom is exactly equal to the number of protons in the nucleus of the atom, so that the overall positive electric charge provided by its protons is exactly balanced by the overall negative charge provided by the electrons orbiting its nucleus.

The unique properties of each of the chemical elements are determined by their number of neutrons, protons, and electrons. Besides determining the properties of a pure chemical element, the neutron, proton, and electron content of its atoms also determines its behavior in relation to other chemical elements. Although each element behaves differently and has different properties from all of the others, the atoms of different elements can combine with one another to form clusters of atoms called molecules. It is this combination of atoms that accounts for the enormous variety of chemical substances that can be found in nature and created by modern technology.

When scientists first tried to describe the physical and chemical properties of the elements and chemical compounds, which are formed by the combination of atoms of different elements, they soon became buried under a mountain of seemingly unconnected facts. Many early scientists recognized the need to organize this information, and they attempted to discover some sort of order or pattern that could simplify what seemed to them an overwhelming array of chemical facts. The solution to the



The Russian chemist Dmitry Mendeleyev created his periodic table (below) in 1869 while preparing a chemistry textbook for his students.

problem was the so-called periodic table of the chemical elements.

The modern periodic table is based primarily on the work of the Russian chemist Dmitry Ivanovich Mendeleyev (1834–1907) and the German physicist Julius Lothar Meyer (1830–1895). Working independently, both of these scientists developed similar periodic tables within a few months of each other in 1869. Mendeleyev, however, is usually given the credit for having developed the periodic table because he managed to publish his work first.

Mendeleyev, who was a professor of chemistry at the University of St. Petersburg, developed the periodic table while preparing a chemistry textbook for his students. As part of this project, he had written down the properties of the elements on cards. While sorting through these cards he noticed that when the elements were arranged in order of their weight, similar chemical properties occurred repeatedly at regular intervals. Mendeleyev used this observation to construct his periodic table. He placed the elements in horizontal rows, according to their weight, with the lightest in each row at the left end of the row and the heaviest at the right, and with one row beneath the other, so that all elements with similar properties fell into vertical columns.

Much of Mendeleyev's success depended on his placing elements with similar properties in the same group, even though

this left occasional gaps in the table. He reasoned, correctly, that the elements that belonged in the gaps had not yet been discovered. The locations of the gaps enabled Mendeleyev to predict with remarkable accuracy the properties of these yet-to-be found elements.

The table that Mendeleyev developed is in many ways similar to the one we use today. One of the main differences is that Mendeleyev's table lacks the column containing the elements helium through radon. In Mendeleyev's time none of the elements in this column had yet been found because they are relatively rare and because they show no tendency to undergo chemical reactions. Occasionally, Mendeleyev was forced to switch the order of his elements to make the table come out right, placing elements with greater atomic weights ahead of those with smaller weights. The atomic weight of an element is the average weight of all the atoms that form the element. It is mostly determined by the total number of protons and neutrons the atoms contain. Electrons are so much lighter than these

1		xviii			PRINCI	PLES (DE CHE	MISTR:	Y	
The latter The		PERI	DDIC S	FSTEM	OF TH	E ELE	MENTS	IN GI	OUPS .	AND SERIES.
1	ries						F ELEM	ENTS		
1	35	0	I	II	III	IV	v	VI	VII	VIII
Record R	ı	-	gen H	-		-	_	-		
A Ar Fotas Fot	2	He.	Li- thium Li 7:03	Beryl- lium Be 9-1	Boron II:0	Car- bon C	Nitro- gen N	Oxy- gen O	rine F	
A A For Fo	3	Neon Ne	So- diam Na 23:05	Mag- nesium M.g 24·3	Alu- minian Al 27-0	Sitt- con Si 28-4	Phos- phorus P 31-6	Sul phun S 32-06	Chlo- rine Cl 35-45	The state of the s
X X Cas Cas Bis Lan Cas Ca	4	Ar- gen Ar	Potas- sium K 39-1	Cal- elom Ca 40-1	Scan- dium Sc 44·1	Tita- nium Ti 48-1	Vans- diam V	Chro- mium Cr 59-1	Man- ganese Mn 55-0	Iron balt Nickel
X X Cas Cas Bas Lan Cas Ca	5		Cop- per Cu 63-6	Zine Zn 65:4	Gal- llum Ga 70-0	Ger- manium Ge 72·3	Ar- senic As 75	Sele- ulum Se 79	Bro- mine Br 79-95	
Steel Case		Kryp- ton Kr 81-8	Rubi- diam Rb 85.4 Sil- ver Ag	Stron- tium Sr 87-6 Cad- mium Cd	Yt- trium Y 89-0 In- dium In	Zirco- niem Zr 90.6 Tiu Sn	Nie- bium Nb 94-0 Anti- mony Sb	Molyb- denum Mo 96:0 Tellu- rium Te	Iodine	Buthe-Rho-Palia nium diam dium Ru Rh Pd 101-7 103-0 106-
10	8	Xenon Xe	Cæ- sium Cs	Ba- rium Ba	Lan- thanum La	rium Ce	1200	127	127	
Radium Thought Uradium thum the Uradium the Uradium thum the Uradium thum the Uradium thum the Uradium the Uradium thum the Uradium the Uradium thum the Uradium thum the Uradium thum the Uradium the Uradium thum the Uradium	9		-	-				_		
Radium Thought Uradium thum the Uradium the Uradium thum the Uradium thum the Uradium thum the Uradium the Uradium thum the Uradium the Uradium thum the Uradium thum the Uradium thum the Uradium the Uradium thum the Uradium	10		_	_	Ytter- bium Yb 173	_	Tan- talum Ta 183	Tung- sten W		os Ir Pt (
Ra- slow Ura- slow unu 12 - Rd - Th - U	11		Au	Mer- cury Hg 200-0	That- lium Tl 204-1	Lead Pb 206-9	Bis- muth Bi 208		_	
224 232 239	12			Ra- dlum Rd		The-		Ura- nium U		

nuclear particles that they contribute very little to the weight of an atom. Apparently, listing elements in order of their atomic weights did not always work. It was not until the beginning of the 20th century, with the knowledge gained about the structure of the atom, that the correct way of ordering the elements was discovered and the present periodic table was formulated.

THE NUCLEAR ATOM

The key event that led to the modern understanding of the atom was the discovery that atoms are made up of electrons, protons, and neutrons. Thus, despite its name, which derives from the Greek word for "indivisible," the atom could indeed be divided into smaller components.

In April 1897, Joseph John Thompson, professor of physics and director of the Cavendish Laboratory at Cambridge University in England, announced the discovery of the electron. Thompson reported that this tiny particle had a negative electric charge and a mass of about one two-thousandth of that of the lightest atom. Thompson's momentous discovery of a particle of matter smaller than the atom so startled his colleagues that many thought he had been "pulling their legs." It was no joke, however, and in Thompson's own words: "The production of electrons essentially involves the splitting up of the atom, [with] a part of the mass of the atom getting free and becoming detached from the original atom—that part being one or more electrons."

Ernest Rutherford, the distinguished New Zealand physicist who had been a pupil of Thompson's and who was a professor of physics at Cambridge University, supplied the next step toward the modern understanding of the atom in 1911. He discovered that the atom had a nucleus and that one of the important particles that occupied the nucleus was the positively charged proton.

As a probe for his study of the atom, Rutherford used the newly discovered phenomenon of radioactivity. Radioactive atoms, like uranium and radium, are unstable, and their nuclei spontaneously disintegrate. One of the products of this disintegration is a massive, positively charged particle called an alpha particle. At the time of its discovery, Rutherford did not know that the alpha particle was the nucleus of a helium atom, consisting of two protons and two neutrons. He therefore used the first letter of the Greek alphabet, *alpha*, to identify this particle and distinguish it from the other products given off by radioactive atoms.

Because the atom was too small to observe directly,

10 The Periodic Table

Rutherford's brilliant idea was to use alpha particles as projectiles, firing them at atoms and observing how they scattered. This was like firing bullets at a sealed box and deducing the contents of the box by seeing how the bullets bounced. His target atoms were at first gold atoms contained in very thin sheets of gold foil. Gold was used because it is possible to make gold foil that is very thin, often thinner than fine paper. Rutherford observed that although most of the alpha particles passed right through the target, many were deflected at very large angles. Some were even deflected backward, as if they had hit a stone wall. Rutherford was so astonished by this that he compared it to "firing a 15-inch shell at a piece of tissue paper and having it come back and hit the gunner." Because most of the alpha particles went right through the foil, he reasoned that the atom was mainly empty space but that it must contain a small, heavy, positively charged core that was capable of repelling and scattering the projectiles fired at it. Rutherford called this massive core the nucleus of the atom.

After Rutherford's discovery of the nucleus, it became obvious that the nucleus of hydrogen, the lightest of the atoms, must play a fundamental role in the structure of all atoms. In 1920, he proposed to call this particle the proton, the name by which it has been known ever since.

Finally, in 1932, the British physicist Sir James Chadwick, who also worked at the Cavendish Laboratory in Cambridge, discovered that yet another particle existed in the nucleus of atoms. This new particle was the neutron. It has a mass close to the mass of the proton, but it has no electric charge.

These fundamental discoveries, coupled with the work of a brilliant young English physicist named Henry Moseley, ultimately led to the reason for Mendeleyev's success with the periodic table. Moseley, just before World War I, had been investigating the X rays given off by various elements. X rays are a very penetrating form of radiation usually produced by accelerating electrons to high speeds and then abruptly stopping them by having them smash into a metal target. The collision causes the target to give off X rays. When different elements are used as targets, the X rays have different properties. Each element has its own set of characteristic X rays. They are almost like a fingerprint of the element. Moseley was able to relate the properties of the X rays to the number of protons contained in the element. He discovered that every element had a different number of protons in the nucleus. The number of protons came to be called the atomic number of the element, represented by the letter Z, and it was always a whole number.

Atoms are normally electrically neutral, with an equal number of electrons and protons. This means, for example, that carbon, with an atomic number of 6, has six protons in its nucleus and six electrons outside the nucleus.

ISOTOPES

Moseley's experiments demonstrated that what distinguishes one element from another is its atomic number, the number of protons in the nucleus of its atoms, not its atomic weight, which is a measure of the total number of protons and neutrons in the nucleus. The correct way of ordering the elements in the periodic table was, therefore, by their atomic number, and not, as Mendeleyev had thought, by their atomic weight.

Although all the atoms of a given element have the same number of protons, they can have different numbers of neutrons. This explains, for example, why there are three different species of the element hydrogen. Ordinarily, a hydrogen atom has a lone proton in its nucleus and no neutrons. A heavier form of hydrogen, called deuterium, also has a single proton in the nucleus but contains a neutron as well. A still heavier form of hydrogen, known as tritium, has two neutrons in addition to the proton. These three species are called isotopes of the element hydrogen. Yet even though a deuterium atom, because of its extra neutron, weighs twice as much as an ordinary hydrogen atom, its chemical behavior is similar to that of hydrogen, indicating that the number of protons in the nucleus is what determines the behavior of each element.

Like hydrogen, the majority of the elements have isotopes. Some elements have only two isotopes, while others can have as many as eight or nine. It is a remarkable fact that the relative percentage, or abundance, of each of the various isotopes of any element is the same all over the Earth.

The existence of isotopes also explains why the atomic weight is an unreliable indicator of the position of an element in the periodic table. For any element, the atomic weight really measures the average weight of a mixture of its different isotopes. On this basis, it is possible for an element like argon, which has an atomic number of 18, to exist in a mixture of isotopes that have a greater average atomic weight than that of potassium, whose atomic number is 19. The atomic weight in the periodic table is often a number with a decimal fraction. The atomic weight of calcium, Ca, for example, is 40.08. The nucleus of the calcium atom cannot